

1. Evaluate.

$$\text{a. } \log 1000 = 3 \quad \text{b. } \log_4 1 = 0 \quad \text{c. } \log_3 27 = 3 \quad \text{d. } \log_2 \left(\frac{1}{4}\right) = -2$$

2. Solve for x .

$$\text{a. } \log_4 x = 2$$

$$4^2 = x \\ x = 16 \\ \{16\}$$

$$\text{b. } \log_{\frac{1}{3}} x = 4$$

$$\left(\frac{1}{3}\right)^4 = x \\ \left\{\frac{1}{81}\right\} = x$$

$$\text{c. } \log(2x + 1) = 2$$

$$10^2 = 2x + 1 \\ 100 = 2x + 1 \\ \frac{99}{2} = 2x \\ \left\{\frac{99}{2}\right\}$$

$$\text{d. } \ln e^{3x-1} = 5$$

$$3x - 1 = 5 \\ 3x = 6 \\ \frac{3x}{3} = \frac{6}{3} \\ x = 2 \\ \{2\}$$

3. Use log properties to solve the equation. $\log_3 x = \log_3 7 + \log_3 3$

$$\log_3 x = \log_3 21$$

$$x = 21 \\ \{21\}$$

4. Use log properties to simplify the expression. $2 \log_{10} 5 + \log_{10} 8 - \log_{10} 2$

$$\log\left(\frac{25 \cdot 8}{2}\right) \\ = \log(100) = 2$$

5. Use log properties to expand as much as possible.

$$\text{a. } \log_2(6x^{-3}y^7)$$

$$\text{b. } \ln \frac{e^y}{\sqrt[3]{x}}$$

$$= \log_2 6 - 3 \log_2 x + 7 \log_2 y$$

$$= \ln e^y - \frac{1}{3} \ln x \\ = y - \frac{1}{3} \ln x$$

$$\text{c. } \log_5 \left(\frac{x^2 - 7x - 18}{5} \right)$$

$$= \log_5(x-9)(x+2) - \log_5 5 \\ = \log_5(x-9) + \log(x+2) - 1$$

6. Solve and write the exact solution(s) in a solution set. Show domain restrictions where appropriate.

$$a. 2^{4-3x} = \frac{1}{128}$$

$$2^{4-3x} = 2^{-7}$$

$$4-3x = -7$$

$$\begin{matrix} -3x \\ -3 \end{matrix} = \begin{matrix} -11 \\ -3 \end{matrix}$$

$$x = \frac{11}{3}$$

$$\left\{ \frac{11}{3} \right\}$$

$$b. \log_4(x+2) + \log_4(x-1) = 1$$

$$\log_4(x+2)(x-1) = 1$$

$$4^1 = x^2 - x + 2x - 2$$

$$0 = x^2 + x - 6$$

$$0 = (x+3)(x-2)$$

$$x = -3, 2$$

$$\{2\}$$

Check domain

$$x+2 > 0 \text{ and } x-1 > 0$$

$$x > -2 \text{ and } x > 1$$

7. LaShonda invests into a corporate bond at 4.5 percent compounded quarterly. If she wants to double her money, how many years will it take? How long will it take to triple her money?

$$A = P \left(1 + \frac{r}{n}\right)^{nt}$$

$$\frac{2P}{P} = \left(1 + \frac{0.045}{4}\right)^{4t}$$

$$2 = \left(1 + \frac{0.045}{4}\right)^{4t}$$

$$\ln 2 = \ln \left(1 + \frac{0.045}{4}\right)^{4t}$$

$$\frac{\ln 2}{4 \ln \left(1 + \frac{0.045}{4}\right)} = \frac{4t \ln \left(1 + \frac{0.045}{4}\right)}{4 \ln \left(1 + \frac{0.045}{4}\right)}$$

$$15.49 \approx t$$

It will take about 15.5 years to double

To triple:

$$\frac{\ln 3}{4 \ln \left(1 + \frac{0.045}{4}\right)} \approx 24.55 \text{ years}$$

8. The half-life of Carbon 14 is 5,715 years. How much of a sample of 100 grams of Carbon 14 will remain 12,000 years after its creation?

$$\textcircled{1} \quad N = N_0 e^{kt}$$

$$\text{Find } k \quad \frac{1}{2} \frac{N_0}{N_0} = \frac{N_0}{N_0} e^{k(5715)}$$

$$\frac{1}{2} = e^{5715k}$$

$$\ln \frac{1}{2} = \ln e^{5715k}$$

$$\frac{\ln \frac{1}{2}}{5715} = \frac{5715k}{5715}$$

$$-.00012 = k$$

$$\textcircled{2} \quad \text{use } N = N_0 e^{-0.00012t}$$

$$N = 100 e^{-0.00012(12000)}$$

$$\approx 23.69$$

About 23.7 grams would be left after 12,000 years.

9. For each of the following functions:

- State the domain of the function
- Describe the end behavior of the function.
- If the function has any asymptotes, state the equation for each asymptote and identify the type of asymptote (vertical, horizontal, or oblique).
- If the function has any holes, state the location of each hole.

a. $f(x) = \frac{2x - 6x^2}{3x - 1}$

$$= \frac{2x(1 - 3x)}{3x - 1}$$

$$= \frac{-2x(3x - 1)}{(3x - 1)}$$

$$= -2x, x \neq \frac{1}{3}$$

Domain: $\{x | x \neq \frac{1}{3}\}$

As $x \rightarrow \infty, y \rightarrow -\infty$

As $x \rightarrow -\infty, y \rightarrow \infty$

No Asymptotes

d. $f(x) = 3x^2 - 1$

Domain: \mathbb{R}

End behavior: $3x^2 \nearrow$

as $x \rightarrow \infty, y \rightarrow \infty$

as $x \rightarrow -\infty, y \rightarrow \infty$

No asymptotes

No holes

b. $f(x) = -2x^2 + 3x - 4$

Domain: \mathbb{R}

End behavior:

$$-2x^2 \searrow$$

as $x \rightarrow \infty, y \rightarrow -\infty$

as $x \rightarrow -\infty, y \rightarrow -\infty$

No holes

No Asymptotes

c. $f(x) = \frac{3x - 1}{3 + x^2}$

$$3 + x^2 \neq 0$$

$$x^2 \neq -3$$

$$x \neq \pm \sqrt{-3}$$

Imaginary numbers

Domain: \mathbb{R}

No vertical asymptote

H.A.: $\frac{3x}{x^2} = \frac{3}{x} \rightarrow 0$

$y = 0$ as $x \rightarrow \pm \infty$

as $x \rightarrow \infty, y \rightarrow 0$

as $x \rightarrow -\infty, y \rightarrow 0$

No holes

e. $f(x) = \frac{(x+2)^2(x-2)(x+4)}{(1-x^2)(x+4)}$

$$= \frac{(x+2)^2(x-2)}{(1-x)(1+x)}, x \neq -4$$

Domain: $\{x | x \neq -4, -1, 1\}$

Vertical asymptotes: $x = -1$
 $x = 1$

Horizontal asymptotes:

$$\frac{x^3}{-x^2} = -x \text{ no H.A.}$$


(oblique asymptote)

as $x \rightarrow \infty, y \rightarrow -\infty$

as $x \rightarrow -\infty, y \rightarrow \infty$

10. Decide whether the function shown is most likely a polynomial, rational or something else. After you have decided, find a formula for the graph, including the coefficient k .

a.



Rational

$$y = \frac{k(x-2)}{(x+2)(x-2)}$$

$$-1 = \frac{k(-3-2)}{(-3+2)(-3-2)}$$

$$\therefore -1 = \frac{k}{-1} \cdot -1$$

$$1 = k$$

$$y = \frac{(x-2)}{(x+2)(x-2)}$$

c. Rational


$$y = \frac{k(x^2+1)}{(x-3)(x+3)}$$

$$k = -4$$

$$y = \frac{-4(x^2+1)}{(x-3)(x+3)}$$

Cara Lee

b.



Polynomial

$$y = k(x+4)^2(x-0)(x-2)(x-5)$$


$$6 = k(-3+4)^2(-3)(-3-2)(-3-5)$$

$$6 = k(1)^2(-3)(-5)(-8)$$

$$\frac{6}{-120} = k \cdot \frac{-120}{-120} \quad k = -\frac{1}{20}$$

$$y = -\frac{1}{20}x^2(x+4)^2(x-2)(x-5)$$

c.



challenge problem
no real zeros

but the degree must be the same on top + bottom to get a H.A. at $y = -4$

11. Draw complete graphs of each function, following the procedures outline in the rational function and polynomial function lecture notes.

a. Graph $y = \frac{(x+2)^2(x-2)(x+4)}{(1-x^2)(x+4)}$

Same as 10e on the previous page:


H.A. = $-x$ oblique
 Challenge problem

Zeros: -2, mult 2
 2, mult 1

Hole at $(-4, 1.6)$

$$y = \frac{(-4+2)^2(-4-2)}{(1-(-4)^2)} \\ = \frac{4(-6)}{1-16} = \frac{-24}{-15}$$

$$y\text{-int} = \frac{(2)^2(-2)(4)}{(1)(4)} = -8$$



(We would need to plot more points for this graph - I used table in the calculator)

b. Graph $y = -\frac{(x-3)^2(x+2)}{3}$

$$y = -\frac{1}{3}(x-3)^2(x+2)$$

zeros	mult
3	2
-2	1

end behavior: $-\frac{1}{3}x^3$

$$y\text{-int}: -\frac{1}{3}(0-3)^2(0+2) \\ = -\frac{1}{3}(2)^2(2) \\ = -\frac{8}{3} \text{ or } -2\frac{2}{3}$$

